

Low T Quiz Results



© TESTOSTERONE UNDECANOATE CAPSULES

The Androgen Deficiency in Aging Males (ADAM) questionnaire on the KYZATREX website collected data from **779** participants over 16 weeks. The results highlight significant challenges faced by men regarding **energy, libido, strength** and overall **life satisfaction**, underscoring the potential impact of testosterone deficiency on quality of life and well-being.

KYZATREX® (testosterone undecanoate) is a prescription drug that is used to treat adult men who have low or no testosterone levels due to certain medical conditions.

KYZATREX is a controlled substance (CIII) because it contains testosterone. It is not known if KYZATREX is safe or effective in males younger than 18 years old. Improper use may affect bone growth in children. KYZATREX is not meant for use by women.

62%

of men rated their **enjoyment of life** as **poor or terrible**.



66%

of men rated their **strength and/or endurance** as **poor or terrible**.




72%

of men rated their **sports ability** over the past 4 weeks as **poor or terrible**.



76%

of men rated their **energy level** as **poor or terrible**.



75% of men rated their **libido** as **poor or terrible**.

46%

of men rated their **happiness level** as **poor or terrible**.



59%

of men **fall asleep**

at least **3-4x/week** after eating dinner.

3-4x/week » 21%
5-6x/week » 17%
Every night » 21%



53%

of men rated their **work performance**

over the past 4 weeks as **poor or terrible**.



72%

of men rated the **strength of their erections**

as **somewhat or extremely weak**.



IMPORTANT SAFETY INFORMATION FOR KYZATREX®

KYZATREX can increase blood pressure, which can increase the risk of having a heart attack or stroke and can increase risk of death due to a heart attack or stroke. Your risk may be greater if you have already had a heart attack or stroke or if you have other risk factors for heart attack or stroke.

- If your blood pressure increases while on KYZATREX, blood pressure medicines may need to be started. If you are currently taking blood pressure medicines, they may need to be changed or new blood pressure medicines may need to be added to control your blood pressure.
- If your blood pressure cannot be controlled, KYZATREX may need to be stopped.
- Your healthcare provider will monitor your blood pressure while you are being treated with

PLEASE SEE ADDITIONAL IMPORTANT SAFETY INFORMATION, INCLUDING BOXED WARNING ON THE NEXT PAGE.

IMPORTANT SAFETY INFORMATION

Use

KYZATREX® (testosterone undecanoate) is a prescription drug that is used to treat adult men who have low or no testosterone levels due to certain medical conditions.

KYZATREX is a controlled substance (CIII) because it contains testosterone. It is not known if KYZATREX is safe or effective in males younger than 18 years old. Improper use may affect bone growth in children. KYZATREX is not meant for use by women.

Important Safety Information For KYZATREX®

KYZATREX can increase blood pressure, which can increase the risk of having a heart attack or stroke and can increase risk of death due to a heart attack or stroke. Your risk may be greater if you have already had a heart attack or stroke or if you have other risk factors for heart attack or stroke.

- If your blood pressure increases while on KYZATREX, blood pressure medicines may need to be started. If you are currently taking blood pressure medicines, they may need to be changed or new blood pressure medicines may need to be added to control your blood pressure.
- If your blood pressure cannot be controlled, KYZATREX may need to be stopped.
- Your healthcare provider will monitor your blood pressure while you are being treated with KYZATREX.

Do not take KYZATREX if you: have breast cancer; have or might have prostate cancer; are a woman who is pregnant (KYZATREX may harm your unborn baby); are allergic to KYZATREX or any of its ingredients; or have low testosterone without certain medical conditions (e.g., do not take KYZATREX if you have low testosterone due to age).

Before you take KYZATREX, tell your healthcare provider about all of your medical conditions, including if you: have high blood pressure or are treated for high blood pressure; have a history of diabetes; have heart problems; have high red blood cell count (hematocrit) or high hemoglobin laboratory value; have urinary problems due to an enlarged prostate; have liver or kidney problems; or have problems breathing while you sleep (sleep apnea).

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Taking KYZATREX with certain other medicines can cause adverse effects. **Especially, tell your healthcare provider if you take:** insulin; medicines that decrease blood clotting (blood thinners); corticosteroids; or medicines that increase blood pressure, such as some cold medicines and pain medicines.

KYZATREX may cause other serious side effects including

- **Increase in red blood cell count (hematocrit) or hemoglobin,** which can increase the risk of blood clots, strokes, and heart attacks. You may need to stop KYZATREX if your red blood cell count increases.
- **If you already have an enlarged prostate, your signs and symptoms may worsen while taking KYZATREX. These may include: increased urination at night; trouble starting your urine stream; urinating many times during the day; urge to go to the bathroom right away; a urine accident; inability to pass urine or weak urine flow.**
- **Increased risk of prostate cancer.**
- **Blood clots in the legs or lungs.** Signs and symptoms of a blood clot in your leg can include pain, swelling or redness. Signs and symptoms of a blood clot

in your lungs can include difficulty breathing or chest pain.

- **Abuse.** Testosterone can be abused when taken at higher than prescribed doses and when used with other anabolic androgenic steroids. Abuse can cause serious heart and psychological side effects.
- **In large doses, KYZATREX may lower your sperm count.**
- **Liver problems.** Symptoms of liver problems may include: nausea or vomiting; yellowing of your skin or whites of your eyes; dark urine; pain on the right side of your stomach area (abdominal pain).
- **Swelling of your ankles, feet, or body (edema), with or without heart failure.**
- **Enlarged or painful breasts.**
- **Breathing problems while you sleep (sleep apnea).**

Call your healthcare provider right away if you have any of the serious side effects listed above.

The most common side effect of KYZATREX is high blood pressure. Other side effects may include: headache, joint or back pain, diarrhea, increased red blood cell count, anxiety, constipation, swelling of the legs, and increased prostate specific antigen (PSA) levels.

These are not all the possible side effects of KYZATREX. For more information, ask your healthcare provider or pharmacist.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088. You may also report side effects to Marius by visiting www.mariuspharma.com.

Keep KYZATREX and all medicines out of the reach of children.

See Full Prescribing Information and Medication Guide for KYZATREX.



Marius Pharmaceuticals
2301 Sugar Bush Road, Suite 510
Raleigh, NC 27612

919-374-1913
mariuspharma.com

KYZATREX® and its associated logo are registered trademarks of Marius Pharmaceuticals.

©2024 Marius Pharmaceuticals. All rights reserved. Rev. 11/2024