

## MEDICATION GUIDE

### KYZATREX® (ky-ZAH-treks)

(testosterone undecanoate) capsules, for oral use, CIII

#### What is KYZATREX®?

- KYZATREX® is a prescription medicine that contains testosterone. KYZATREX® is used to treat adult men who have low or no testosterone due to certain medical conditions.
- It is not known if KYZATREX® is safe or effective in children younger than 18 years old. Improper use of KYZATREX® may affect bone growth in children.
- KYZATREX® is a controlled substance (CIII) because it contains testosterone that can be a target for people who abuse prescription medicines. Keep your KYZATREX® in a safe place to protect it. Never give your KYZATREX® to anyone else, even if they have the same symptoms you have. Selling or giving away this medicine may harm others and is against the law.
- KYZATREX® is not meant for use by women.

#### Do not take KYZATREX® if you:

- have breast cancer.
- have or might have prostate cancer.
- are a woman who is pregnant. KYZATREX may harm your unborn baby.
- are allergic to KYZATREX® or any ingredients in KYZATREX®. See the end of this Medication Guide for a complete list of ingredients in KYZATREX®.

#### Before you take KYZATREX®, tell your healthcare provider about all of your medical conditions, including if you:

- have high blood pressure or are being treated for high blood pressure.
- have a history of diabetes.
- have heart problems.
- have high red blood cell count (hematocrit) or high hemoglobin laboratory value.
- have urinary problems due to an enlarged prostate.
- have liver or kidney problems.
- have problems breathing while you sleep (sleep apnea).

**Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.** Taking KYZATREX® with certain other medicines can affect each other.

#### Especially, tell your healthcare provider if you take:

- insulin
- medicines that decrease blood clotting (blood thinners)
- corticosteroids
- medicines that increase blood pressure such as some cold medicine and pain medicines

Know the medicines you take. Ask your healthcare provider or pharmacist for a list of these medicines, if you are not sure. Keep a list of them and show it to your healthcare provider and pharmacist when you get a new medicine.

#### How should I take KYZATREX®?

- Take KYZATREX® exactly as your healthcare provider tells you take it.
- Take KYZATREX® by mouth 2 times daily. Take 1 time in the morning and take 1 time in the evening. If your dose is 100 mg daily, take 1 time in the morning.
- Take KYZATREX® with food.
- Your healthcare provider may change your KYZATREX® dose. Do not change your KYZATREX® dose without talking to your healthcare provider.

#### What are the possible side effects of KYZATREX®?

**KYZATREX® may cause serious side effects including:**

- See “**What is the most important information I should know about KYZATREX®?**”
- **Increase in red blood cell count (hematocrit) or hemoglobin.**
  - KYZATREX® increases red blood cell counts in some patients. High red blood cell counts increase the risk of blood clots, strokes, and heart attacks.
  - You may need to stop KYZATREX if your red blood cell count increases.
  - Your healthcare provider should check your red blood cell count and hemoglobin while you take KYZATREX®.
- **If you already have an enlarged prostate, your signs and symptoms may worsen while taking KYZATREX®.** These may include:
  - increased urination at night
  - trouble starting your urine stream
  - urinating many times during the day
  - urge to go to the bathroom right away
  - a urine accident
  - inability to pass urine or weak urine flow
- **Increased risk of prostate cancer.** Your healthcare provider should check you for prostate cancer or any other prostate problems before you start and while you take KYZATREX®
- **Blood clots in the legs or lungs.** Signs and symptoms of a blood clot in your leg can include pain, swelling or redness. Signs and symptoms of a blood clot in your lungs can include difficulty breathing or chest pain.
- **Increase in blood pressure.** KYZATREX can increase your blood pressure. Increases in blood pressure can increase the risk of heart attack or stroke over time. If your blood pressure increases while on KYZATREX, blood pressure medicines may need to be started. If you are taking blood pressure medicines, new blood pressure medicines may need to be added or your current blood pressure medicines may need to be adjusted to control your blood pressure. If your blood pressure cannot be controlled, KYZATREX may need to be stopped. Your healthcare provider will monitor your blood pressure while you are being treated with KYZATREX.
- **Abuse.** Testosterone can be abused, when taken at higher than prescribed doses and when used with other anabolic androgenic steroids. Abuse can cause serious heart and psychological side effects. Your healthcare provider should check you for signs of abuse before and during treatment with KYZATREX®.
- **In large doses KYZATREX® may lower your sperm count.**
- **Liver problems.** Symptoms of liver problems may include:
  - nausea or vomiting
  - yellowing of your skin or whites of your eyes
  - dark urine
  - pain on the right side of your stomach area (abdominal pain)
- **Swelling of your ankles, feet, or body (edema), with or without heart failure.**
- **Enlarged or painful breasts.**
- **Breathing problems while you sleep (sleep apnea).**

**Call your healthcare provider right away if you have any of the serious side effects listed above.**

**The most common side effect of KYZATREX® is high blood pressure.**

**Other side effects may include** headache, joint or back pain, diarrhea, increased red blood cell count, anxiety, constipation, swelling of the legs, and increased prostate specific antigen (PSA) levels.

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects of KYZATREX®. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

#### **How should I store KYZATREX®?**

- Store KYZATREX® at room temperature between 68°F to 77°F (20°C to 25°C).
- Store KYZATREX® in a dry place.

**Keep KYZATREX and all medicines out of the reach of children.**

#### **How should I throw away (dispose of) KYZATREX®?**

- Throw away unused KYZATREX® via a take-back option.
- If a take-back option is unavailable, follow FDA instructions at [www.fda.gov/drugdisposal](http://www.fda.gov/drugdisposal) for properly throwing away medicine.

#### **General information about the safe and effective use of KYZATREX®**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use KYZATREX® for a condition for which it was not prescribed. Do not give KYZATREX® to other people, even if they have the same symptoms you have. It may harm them. You can ask your pharmacist or healthcare provider for information about KYZATREX® that is written for health professionals.

**What are the ingredients in KYZATREX®?**

**Active ingredient:** testosterone undecanoate

**Inactive ingredients:** DL-alpha-tocopheryl acetate (Vitamin E), phytosterol esters, polyoxyl 40 hydrogenated castor oil and propylene glycol monolaurate. The ingredients of the gelatin capsule shells are gelatin, glycerin, purified water, sorbitol, and titanium dioxide.

Marketed by:

Marius Pharmaceuticals  
Raleigh, NC 27612

For more information, go to [www.KYZATREX.com](http://www.KYZATREX.com) or call 1-833-949-5040

This Medication Guide has been approved by the U.S. Food and Drug Administration

Issued: 07/2025